

## Claudia Bogado

Contact Information	<ul> <li>Av Medrano 134 floor 6 Caba. Bs As. Argentina. Zip Code C1179AAB.</li> <li>Phone 054 011 4983 1589.</li> <li>Email: claudiabogado@imoba.com.ar</li> </ul>
Education	<ul> <li>Dispositión 6677/10 ANMAT, Imoba 2018</li> </ul>
	Good Clinical Practice, Eli Lilly jun 2018.
	Good clinical Practice, The Global Healt Network Oct 2021.
	<ul> <li>Diploma in Good Practices in Biomedical Research. University of San Pablo.Tucumán. August 2020 to July 2021 (238 hours).</li> </ul>
Current employment	Investigaciones Médicas IMOBA SRL
information	<ul> <li>Av. Medrano 134 floor 6 CABA</li> </ul>
	<ul> <li>Study coordinator from 2018 to present.</li> </ul>
<i>Clinical studies in the last 5 year</i>	I8B-MC-ITRN Prospective, double-blind, randomized study of LY900014 compared with insulin Lispro in combination with insulin Glargine or insulin Degludec in adults with type 2 diabetes PRONTO-T2D. From 2018 to 2019. Phase III. Eli Lilly. Studies coordinator.
	I8B-MC-ITRM Prospective, double-blind, randomized study of LY900014 compared with Lispro insulin with an open-label treatment group with postprandial LY900014 in combination with insulin Glargine or insulin Degludec in adults with type 1 diabetes PRONTO-T1D. From 2018 to 2019. Phase III. Eli

Lilly. Studies coordinator.

- CLCZ696D2301 Multicenter, randomized, double-blind, parallel-group, activecontrolled study to evaluate the efficacy and safety of LCZ696, compared to valsartan, on morbidity and mortality in participants with heart failure (Class II-IV of NYHA) and preserved ejection fraction. From 2018 to the present. Phase 3. Novartis. Studies coordinator.
- 1002-043 (CLEAR), entitled "Randomized, double-blind, placebo-controlled study to evaluate the effects of bempedoic acid (ETC-1002) on the occurrence of major cardiovascular events in patients with cardiovascular disease or at high risk of developing it, who do not tolerate statins
- Protocol H9X-MC-GBGL "A randomized, double-blind, parallel-arm study to study the efficacy and safety of investigational doses of Dulaglutide when added to Metformin in patients with type 2 diabetes mellitus." From 2018 to December 2019. Phase III. Coordinator of clinical studies. Eli Lilly.
- K-877-302 Prominent Pemafibrate to reduce cardiovascular outcomes by lowering triglycerides in patients with diabetes. From 2018 to the present. Phase III. Quintiles. Studies coordinator.
- I8F-MC-GPGM Efficacy and safety of Ly3298176 once a week vs insulin glargine in patients with type 2 diabetes and high cardiovascular risk. From March 2018 to July 2021. Phase III. Eli Lilly. Study coordinator.
- I8F-MC-GPGH: Phase 3 open-label, randomized study comparing the effect of LY3298176 vs titrated insulin Degludec on the glycemic control of pts. with type 2 diabetes. Eli Lilly. From July 2018 to April 2021. Phase III. Eli Lilly. Study coordinator.
- I8F-MC-GPGL: Randomized, open-label, phase 3 trial, comparing the efficiency and safety of Tirzepatide vs Semaglutide once a week as an add-on therapy to metformin in patients with type 2 DBT. Eli Lilly. From July 2019 to May 2021. Phase III. Eli Lilly. Study coordinator.
- FINEART s-HF-(Bayer) Multicenter, randomized, double-blind, placebocontrolled parallel group study to evaluate the efficacy and safety of finerenone in morbidity and mortality in patients with Heart Failure (NYHA II-IV) and fraction left ventricular ejection >=40% (LVEF >=40%). Phase 3. Study

Coordinator. From Oct 2020 to present.
I8H-MC-BDCL A Phase 2 parallel, camparator- Controlled Trial to Evaluate the Safefty and Efficacy of LY3209590 in Insulin- Native with Type 2 Diabetes Mellitus. Eli Lilly. Study coordinator.From Nov 2020 to present.
I8F-MC-GPHM Efficacy and SAfefty of Tirzepatide Once Weekly vs Placebo after an intensive Lifestyle Program participants without Type 2 Diabetes who have Obesiy or are Overweight with weight- related Comorbidities: Randomized Double Blind, Placebo Controlled Trial. Study coordinator. From April 2021 to present.
I8F-MC-GPHD Arandomized, phase 3, open label Trial comparing the effect of the addition of tirzepatide once weekly vs insulin lispro (U100)Three times daily in participants with type 2 diabetes inadequately controlled on insulin glargine (U100) with or without metformin. Study coordinator. From Dec 2020 to present.

Signature:

Date: